Discrete Mathematics I, Sample midterm test Spring 2024

NAME:

NEPTUN code:

Class teacher:

Duration: 90 minutes.

Total score: The maximal total score achievable in the midterm test is 40 points.

Success criterium: In order to successfully past this test, you need to achieve at least 16 points.

Equipment: Only blank papers and pen allowed, no calculators.

Instructions:

Each task must be solved on paper using pen. Please

- write your name, Neptun code and the name of your practice class teacher on the top of this exam paper;
- write your name and Neptun code on the top of each paper you submit;
- note that in most questions **justification is required**. Just an answer to these questions without any proof is worth very few marks only. **Please do not forget to justify your answers**; (Applying and showing the steps of a method learnt in the class where relevant is regarded as sufficient justification.)
- Please note that a 'yes' or no' answer on its own without any justification is worth 0 marks only.
- after finishing the test, place all the papers with your solutions behind this exam paper and in order for the papers to stay together please fold them into half (parallel to the longer side);
- note that the actual exam paper will be shared on Canvas, hence you will be able to obtain it.

In Question 6 you have a choice between two questions: You do not need to solve both of them, please choose just **one** of them.

Thank you and all the best for the test!

Values of some trigonometric functions that may be needed for some questions

| x | 0 | $\frac{\pi}{6}$ | $\frac{\pi}{4}$ | $\frac{\pi}{3}$ | $\frac{\pi}{2}$ | $\frac{2\pi}{3}$ | $\frac{3\pi}{4}$ | $\frac{5\pi}{6}$ | π |
|----------|---|----------------------|---|----------------------|-----------------|----------------------|---|-----------------------|----|
| $\cos x$ | 1 | $\frac{\sqrt{3}}{2}$ | $\frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$ | $\frac{1}{2}$ | 0 | $-\frac{1}{2}$ | $-\frac{\sqrt{2}}{2} = -\frac{1}{\sqrt{2}}$ | $-\frac{\sqrt{3}}{2}$ | -1 |
| $\sin x$ | 0 | $\frac{1}{2}$ | $\frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$ | $\frac{\sqrt{3}}{2}$ | 1 | $\frac{\sqrt{3}}{2}$ | $\frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$ | $\frac{1}{2}$ | 0 |

Questions

- 1. For each of the equalities below decide if it is true for every set A, B and C. Prove your answers.
 - (a) $A \setminus (B \cup C) = (A \setminus B) \cup (A \setminus C).$
 - (b) $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C).$

(4 marks)

- 2. Consider the binary relations $R = \{(1, 4), (1, 5), (2, 7), (3, 2), (6, 1), (6, 6), (7, 4), (7, 5), (7, 7)\}$ and $S = \{(1, 4), (2, 1), (3, 7), (4, 1), (4, 6), (5, 3), (5, 4), (5, 5), (6, 6)\}$. Find each of the following:
 - (a) $dmn(S) \bigtriangleup rng(R)$,
 - (b) $R(\{0, 1, 4, 5, 6, 8\}),$
 - (c) $R^{-1} \circ S|_{\{2,4,5,7\}}$ and

(d) $S \circ R$.

- 3. (a) Let $\rho = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} \mid 2x + 1 = 3y 5\}$. Find $\rho(\{-1, 0, 1, 2, 3, 4\})$ and $\rho^{-1}(\{-1, 0, 1, 2, 3, 4\})$. (b) Given $R = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid 6x - 1 = \frac{4y - 3}{2}\}$ and $S = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid y^2 + 2 = 7x\}$ find the composition $S \circ R$. (7 marks)
- 4. (a) Consider the binary relation $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 4), (3, 1), (3, 3), (4, 2)\}$ on set $X = \{1, 2, 3, 4\}$. Decide, whether R is reflexive, symmetric, transitive and/or anti-symmetric. Justify each of your answers.
 - (b) On set $A = \{1, 2, 3, 4, 5\}$ construct a relation R which satisfies all of the following properties: it is not transitive, not symmetric, not anti-symmetric and $R(\{4\}) = \{1, 5\}$. (6 marks)
- 5. (a) For each of the following relations decide if it is an equivalence relation. Prove your answers. i. $P_{i} \subset \mathbb{P} \times \mathbb{P}$ $= f(x, y) \subset \mathbb{P} \times \mathbb{P}$ $= m \ge 0$

1.
$$R_1 \subseteq \mathbb{R} \times \mathbb{R}, R_1 = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid xy \ge 0\}$$

- ii. $R_2 \subseteq \mathbb{Z} \times \mathbb{Z}, R_2 = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} \mid 4 \mid x y\}$
- (b) For each of those relations above which are equivalence relations, find the partition determined by the equivalence relation. (7 marks)

There are two versions of Question 6, which you can choose from. Please, choose **one** of the two versions. (Even if you solve both of them only one of the two will count.)

Version 1:

- 6. (a) For each of the following relations decide if it is a function.
 - $f_1 \subseteq \mathbb{R} \times \mathbb{R}, f_1 = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid 2x 3 = y^2\}$ $f_2 \subseteq \mathbb{R} \times \mathbb{R}, f_2 = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid 2\cos y = x\}$ $f_3 \subseteq \mathbb{R} \times \mathbb{R}, f_3 = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid y x^2 = 5\}$
 - (b) For each of the above relations which are functions, decide if it is injective, surjective and/or bijective. (7 marks)

Version 2:

6. Using the polar form of complex numbers, calculate the value of:

$$z = \frac{\left(-\frac{7\sqrt{3}}{2} + \frac{7}{2}i\right)^{11}}{\left(-\frac{7}{2} + \frac{7\sqrt{3}}{2}i\right)^{12}}$$

giving your answer in polar form. Find all complex numbers w such that $w^4 = z$, giving your answer in polar form. (7 marks)